International Journal of Nursing and Health Research
www.nursingjournal.in
ISSN: 2664-6420
Received: 12-09-2026, Accepted: 12-10-2026, Published: 27-10-2026
Volume 8, Issue 4, 2026, Page No. 26-32

A critical analysis of mental health nursing interventions in the management of schizophrenia

Rajender Prasad Bhawaria[1], Varsha Arora[1], Rakesh Kumar Yadav[3], Rajni Sharma[3]
[1] Principal, Symbiosis Institute of Nursing, Jaipur, Rajasthan, India
[2] Vice Principal, Symbiosis Institute of Nursing, Jaipur, Rajasthan, India
[3] Assistant Professor, Symbiosis Institute of Nursing, Jaipur, Rajasthan, India
DOI: https://doi.org/10.66856/ijnhr.2026.8.4.8079

Abstract
Schizophrenia is a complex and persistent mental health disorder that can influence a person's thoughts, emotions, behaviour, communication, and ability to carry out everyday activities. While medication remains an important part of treatment, effective care also depends on regular nursing support and appropriate psychosocial interventions. This paper critically examines the role of mental health nursing interventions in the management of schizophrenia, particularly in relation to symptom control, patient safety, treatment adherence, recovery, and quality of life. Important nursing interventions include therapeutic communication, mental health assessment, and medication support, psych education, behavioural management, relapse prevention, family participation, and psychosocial rehabilitation. Developing a trusting relationship with the patient enables nurses to understand individual needs and provide appropriate support. Continuous assessment also helps in recognising changes in mental status, behaviour, and treatment response at an early stage. Nurses contribute to safe medication management by encouraging adherence and observing possible adverse effects. Psych education and family involvement can improve awareness of the illness and help caregivers identify early signs of relapse. Rehabilitation-focused interventions can further assist patients in developing greater independence, social skills, and participation in community life. However, nursing care may be challenged by treatment non-adherence, stigma, cognitive difficulties, inadequate family support, and limited access to community-based mental health services. Conclusively, effective mental health nursing requires a person-centred and recovery-focused approach that combines clinical care with emotional, social, and family support.

Keywords: Schizophrenia, mental health nursing, therapeutic communication, psych education

Introduction

Schizophrenia is a severe and long-term mental health condition that can influence a person's thoughts, perception, emotions, behaviour, relationships, and ability to perform daily activities. The World Health Organization (WHO) estimates that around 23 million people worldwide live with schizophrenia, which is approximately 1 in every 345 people. The condition can have a major effect on education, employment, family life, social relationships, and independent living. WHO also reports that more than two-thirds of people experiencing psychosis worldwide do not receive specialist mental health care? [1] This treatment gap shows that schizophrenia is not only a clinical problem but also a continuing public health and social-care concern. Its management therefore requires regular medical treatment along with psychological, social, rehabilitative, and nursing support.

Treatment of schizophrenia generally includes antipsychotic medicines, psychological interventions, psych education, family support, rehabilitation, and regular follow-up. Mental health nurses have an important role within this multidisciplinary system because they often spend considerable time with patients and are closely involved in their day-to-day care. Their responsibilities include assessing mental health status, establishing therapeutic communication, administering and monitoring medicines, maintaining patient safety, managing disturbed behaviour, providing health education, supporting families, and identifying early signs of relapse. In clinical practice, nurses may notice changes in sleep, communication, self-care, behaviour, medication use, or psychotic symptoms before these changes become more severe [2]. Early recognition can allow timely intervention and may help prevent further deterioration. Effective nursing care can therefore support symptom management while also promoting treatment continuity, safety, recovery, and quality of life.

The importance of mental health nursing is also recognised within India's mental health service system. The National Mental Health Programme (NMHP) was introduced in 1982 to improve access to mental healthcare, integrate mental health services with general healthcare, promote community participation, and develop trained human resources. The District Mental Health Programme (DMHP) was introduced in 1996 to strengthen mental health services at the district level through early identification and treatment, training, public awareness, and community-based follow-up. According to current government information, DMHP has been approved for implementation in 767 districts, with multidisciplinary teams that include psychiatric or community nurses [3]. This structure highlights the importance of nursing professionals in providing mental healthcare outside specialised psychiatric hospitals.

India has also expanded the use of technology through the National Tele Mental Health Programme, including Tele-MANAS, which provides round-the-clock access to mental health support. The service is designed to provide counselling, psychosocial support, medical advice, specialist consultation, follow-up, and referral to appropriate face-to-face services when required [4]. Such services have practical importance for people with long-term mental health

conditions because continued contact and follow-up can help maintain treatment and identify difficulties at an earlier stage. For individuals with schizophrenia, community and tele-mental health services may also help reduce gaps in care, particularly when regular access to specialist services is difficult.

The National Mental Health Policy 2014 further provides a policy framework based on principles such as equity, human rights, participation, recovery, social inclusion, and access to appropriate mental healthcare [5]. These principles are directly relevant to mental health nursing. Nursing care should not be limited to observing symptoms and giving medicines. Nurses also need to protect patient dignity, encourage involvement in care decisions, provide understandable information, involve families appropriately, address stigma, and support patients in developing greater independence [6]. This broader role reflects the shift from a purely illness-focused model toward a more person-centred approach to mental healthcare.

However, the presence of nursing interventions does not guarantee the same level of benefit for every individual with schizophrenia. Treatment adherence can be affected by medication side effects, limited insight, cognitive problems, stigma, financial difficulties, family circumstances, and lack of regular community support. Similarly, concentrating only on reducing hallucinations, delusions, or disturbed behaviour may leave other important areas of recovery insufficiently addressed. Independence, social relationships, self-care, meaningful activity, employment, and quality of life are also important outcomes of long-term mental health care [7]. For this reason, nursing interventions need to be critically examined in terms of their practical value, limitations, and ability to meet individual patient needs.

In this context, the present paper critically analyses mental health nursing interventions used in the management of schizophrenia. It focuses on therapeutic communication, mental health assessment, medication-related care, behavioural support, psych education, family involvement, relapse prevention, and psychosocial rehabilitation. The discussion also considers practical challenges and the relevance of Indian mental health programmes and policies. Particular attention is given to the movement from hospital-based symptom management toward person-centred, recovery-oriented, and community-based care. A balanced approach to nursing can help individuals with schizophrenia achieve greater clinical stability while also supporting dignity, independence, social participation, and long-term recovery.




Fig 1: Clinical Features, Psychosocial Impact, and Nursing Implications in Schizophrenia

Figure 1 presents a simple conceptual view of the main clinical features of schizophrenia, their effects on the individual and family, and the areas in which mental health nursing can provide support. The clinical features are broadly grouped into positive, negative, and cognitive symptoms. Positive symptoms, such as hallucinations, delusions, disorganised speech, and disturbed behaviour, can change how a person understands and responds to the surrounding environment. Negative symptoms, such as low motivation, reduced emotional expression, loss of interest, and withdrawal from others, may gradually affect self-care, education, work, and social activities. Cognitive problems related to attention, memory, concentration, planning, and decision-making can make it difficult for individuals to understand information, follow instructions, and manage their treatment effectively.

The figure also shows that the effects of schizophrenia are not limited to psychiatric symptoms. The condition may create difficulties in daily functioning, independence, relationships, and community participation. Family members and caregivers may also face emotional, social, and financial pressure when providing on-going support with treatment, appointments, personal care, and changes in behaviour. In addition, stigma and negative attitudes toward mental illness can lead to discrimination, social isolation, reduced confidence, and reluctance to seek or continue treatment.

The final part of the figure highlights the practical role of mental health nursing in responding to these needs. Nursing care involves regular assessment, therapeutic communication, medication support, patient and family education, safety management, relapse prevention, and

rehabilitation. These interventions should be adapted to the individual's symptoms, abilities, needs, and social circumstances. The overall purpose is not simply to reduce psychiatric symptoms but to help the person achieve better functioning, greater independence, dignity, social involvement, and sustained recovery.

Mental Health Nursing Interventions in the Management of Schizophrenia

Mental health nurses have an important role in the care of people with schizophrenia because support is often needed over a long period. Nursing care involves much more than giving prescribed medicines. It includes observing the patient's mental and physical condition, building a therapeutic relationship, maintaining safety, providing health education, supporting families, preventing relapse, and helping patients develop skills for daily living. The overall purpose of these interventions is to improve symptom management while also supporting treatment adherence, functioning, independence, and recovery.

Assessment and Continuous Monitoring

Care begins with a careful and regular assessment of the patient. Nurses monitor changes in thoughts, mood, perception, communication, behaviour, sleep, appetite, personal hygiene, and interaction with others. Mental status assessment can help identify hallucinations, delusions, disorganised thinking, confusion, agitation, and other changes in mental functioning. Attention should also be given to self-care and everyday activities because a decline in these areas may occur even when obvious psychotic symptoms are not present.

Regular observation becomes especially important during acute phases of illness. Increased agitation, fear, suspiciousness, aggression, severe withdrawal, or unusual behaviour may indicate worsening symptoms or a possible safety concern. Nurses should record important findings and share significant changes with the treatment team [8]. This allows the patient's care plan and treatment to be reviewed when required.

Therapeutic Communication

Developing a trusting relationship is a central part of mental health nursing. A person experiencing hallucinations, suspiciousness, or delusional ideas may find it difficult to trust others. Nurses should therefore use a calm, respectful, and patient approach. Communication should be clear and simple, particularly when the patient has difficulty concentrating or processing information [9].

Nurses should avoid directly arguing with a patient about a delusional belief or dismissing the person's experience. Instead, they can recognise the emotions associated with the experience and provide reassurance without confirming the false belief. For example, when a patient reports hearing voices, the nurse can acknowledge that the experience is frightening or disturbing and focus on ways of helping the patient feel safe. Consistent and respectful communication can improve cooperation and create a supportive therapeutic environment.

Medication Management and Treatment Adherence

Medication is an important part of schizophrenia treatment, and nurses are closely involved in safe medication management. Their responsibilities include giving medicines according to the prescription, observing treatment response, monitoring possible adverse effects, and explaining the purpose of treatment to the patient. Nurses can also identify difficulties that may lead to missed doses or discontinuation of medication.

Treatment may be interrupted because of unpleasant side effects, lack of understanding, forgetfulness, financial difficulties, or limited awareness of the illness. Instead of treating non-adherence simply as a failure of the patient, nurses should try to understand the underlying reason [10]. This information can then be discussed with the treating team so that appropriate solutions can be considered. Patients should receive information in understandable language and should be involved in treatment discussions as much as possible.

Management of Behavioural Problems and Patient Safety

Some patients may experience severe anxiety, agitation, confusion, or disturbed behaviour during acute psychotic episodes. Protecting the patient and others therefore becomes an important nursing responsibility. Nurses can reduce unnecessary noise and stimulation, identify factors that may increase distress, maintain appropriate observation, and use a calm approach during difficult situations.

Safety assessment should include relevant risks such as severe agitation, self-neglect, suicidal thoughts, threats, previous harmful behaviour, or inability to meet basic needs. When restrictive measures are considered, they should only be used when necessary and in accordance with professional standards, legal requirements, and institutional policies. Wherever possible, nurses should first use supportive and less restrictive approaches while maintaining patient dignity and safety [11].

Psycho-education

Psycho-education helps patients develop a better understanding of schizophrenia and its management. Nurses can explain the nature of the illness, common symptoms, treatment options, medication, possible side effects, relapse warning signs, and the importance of follow-up. Information about sleep, stress management, substance use, and healthy daily routines may also be useful.

Education should match the patient's level of understanding and cognitive ability. Important information may need to be repeated over time rather than provided during a single session. Nurses should also ask patients to explain the information in their own words when appropriate [12]. This helps determine whether the patient has understood the advice and identifies areas that need further explanation.

Family Education and Support

Family members often play an important role in the long-term care of individuals with schizophrenia. They may help with medication, appointments, daily activities, and observation of changes in behaviour. However, families may find these responsibilities difficult when they have limited knowledge about the condition.

Nurses can provide families with practical information about symptoms, treatment, medication, warning signs of relapse, communication, and available support services. Family participation should be guided by the patient's wishes, consent, confidentiality requirements, and clinical circumstances. Nurses should also recognise the needs of

caregivers because prolonged caregiving may cause emotional, social, and financial stress [13].

Relapse Prevention

Because schizophrenia can involve repeated episodes of worsening symptoms, preventing relapse is an important part of continuing care. Nurses can work with patients and families to identify changes that commonly appear before a relapse. These may include disturbed sleep, increasing suspiciousness, and withdrawal, changes in self-care, unusual communication, increased anxiety, or problems with medication.

A practical relapse-prevention plan can record the patient's early warning signs, appropriate actions, healthcare contacts, medication arrangements, and follow-up schedule. When changes are identified early, the treatment team may be able to respond before symptoms become severe [14]. This approach can support continuity of treatment and may help avoid unnecessary emergency care or hospital admission.

Psycho-social Rehabilitation and Functional Support

Recovery should not be judged only by whether psychotic symptoms have decreased. Many people with schizophrenia also need support to improve self-care, communication, social interaction, education, employment, and independent living. Nurses can encourage participation in suitable rehabilitation activities based on the person's abilities and goals.

Goals should be realistic and developed according to individual needs. One person may need help with personal hygiene and medication routines, while another may be ready to work on social skills, vocational activities, or community participation. Small and achievable improvements can gradually increase confidence and independence [15].

Community Follow-Up and Continuity of Care

Schizophrenia often requires continuing care after discharge from hospital. Community-based services can help patients maintain contact with healthcare professionals and receive support before problems become serious. Nurses may contribute through follow-up, medication monitoring, health education, family guidance, and coordination with other professionals.

Continuity becomes especially important when a patient moves between hospital, outpatient, and community services. Poor communication between services can lead to missed appointments, medication difficulties, or delays in responding to worsening symptoms. Good coordination and clear communication between healthcare providers can provide a more stable and connected care pathway [16].

Person-Centred and Recovery-Oriented Care

Mental health nursing should consider the person behind the diagnosis. Person-centred care takes account of the patient's preferences, strengths, abilities, needs, social situation, and personal goals. Wherever possible, patients should be encouraged to participate in decisions about their treatment and daily care.

Recovery does not necessarily mean that every symptom disappears completely. For many individuals, recovery may involve developing confidence, maintaining relationships, improving self-care, participating in education or employment, and living more independently. Nursing care should therefore combine symptom management with support for dignity, autonomy, social participation, and quality of life. A recovery-oriented approach can help patients develop practical strengths and make meaningful progress in their everyday lives [17].

Critical Analysis, Challenges, and Future Directions

Mental health nursing plays an important role in the long-term care of people with schizophrenia. Nursing interventions can help in managing symptoms, improving treatment adherence, maintaining safety, supporting daily activities, and promoting recovery. However, the effectiveness of nursing care is not the same for every patient. Factors such as the severity of symptoms, cognitive difficulties, family support, personal preferences, social circumstances, and availability of mental health services can influence treatment outcomes. Therefore, nursing interventions need to be examined not only in terms of their benefits but also in relation to their practical limitations and the conditions in which they are provided.

Critical Analysis of Nursing Interventions

Assessment and monitoring form the foundation of nursing care in schizophrenia. Regular assessment helps nurses recognise changes in mental status, behaviour, self-care, and response to treatment. One of its major advantages is that early changes in the patient's condition can be identified and addressed before they become more serious. However, assessment can become too focused on psychiatric symptoms such as hallucinations and delusions. This may result in less attention being given to everyday functioning and social needs. A comprehensive assessment should therefore also consider self-care, relationships, treatment adherence, quality of life, and the concerns expressed by the patient.

Therapeutic communication is another important nursing intervention. A calm, respectful, and consistent approach can help patients feel safer and gradually develop trust with the nurse. This trust can improve communication and cooperation with treatment. However, establishing a therapeutic relationship can be difficult when a patient is highly suspicious, frightened, disorganised, or experiencing severe psychotic symptoms. Nurses therefore need patience, appropriate communication skills, and enough time to build a meaningful relationship. They should listen to the patient's experiences without arguing about delusional beliefs or confirming beliefs that are not based in reality.

Medication-related care is also a major responsibility of mental health nurses. Nurses support medication use by providing information, observing treatment response, identifying adverse effects, and discussing difficulties with taking medicines. However, medication adherence should not be viewed simply as the patient's responsibility. Missed medication may result from unpleasant side effects, financial difficulties, limited access to medicines, forgetfulness, lack of understanding, or limited family support. For this reason, medication care should be combined with patient education, discussion of treatment choices, and regular follow-up.

Psycho education and family support can improve long-term management of schizophrenia. Providing clear information about the illness, treatment, medication, and warning signs can help patients and families respond more effectively to difficulties. At the same time, family involvement cannot be

approached in the same way for everyone. Some patients may have little family support, strained relationships, or concerns about privacy. Nursing care should therefore respect the patient's preferences and confidentiality while also identifying alternative sources of support when needed.

Psychosocial rehabilitation is important because controlling psychiatric symptoms does not automatically restore a person's independence. Patients may continue to require help with self-care, communication, social interaction, education, employment, or community participation. Rehabilitation can help individuals gradually develop these abilities and become more independent. However, access to rehabilitation and community-based services is not equal in all settings. Where such services are limited, patients may remain dependent on hospital-based care even when their acute symptoms have improved.

Challenges in Mental Health Nursing Care

Several practical and social factors can reduce the effectiveness of nursing interventions in schizophrenia.

Treatment non-adherence is a significant challenge. Patients may stop or irregularly take medication because of side effects, lack of insight, forgetfulness, stigma, financial problems, or difficulty reaching healthcare services. Simply describing such patients as uncooperative does not address the actual problem. Nurses should try to understand the reasons behind missed treatment and work with the patient and healthcare team to find practical solutions.

Stigma and discrimination can further complicate recovery. Fear of being judged may prevent individuals from discussing their symptoms, seeking help, maintaining relationships, or participating in community activities. Nurses can contribute to reducing stigma by providing respectful care, correcting misconceptions, educating patients and families, and encouraging social inclusion.

Cognitive difficulties may also interfere with treatment. Problems with attention, memory, concentration, and decision-making can make it difficult for patients to remember instructions or follow treatment routines. Therefore, information should be presented in simple language and repeated when required. Teaching should be adjusted according to the individual’s level of understanding rather than using the same approach for every patient.

Family and caregiver burden is another important concern. Families may have to provide on-going assistance with medication, appointments, daily activities, and management of behavioural changes. This can create emotional, social, and financial pressure. If caregivers do not receive adequate support, their ability to continue providing care may be affected. Nurses can help by providing education, emotional support, counselling, and information about available community and support services.

A further challenge is the limited and unequal availability of mental health services and trained professionals. Shortage of staff, heavy workloads, inadequate community facilities, and weak coordination between different healthcare services can limit the amount of individual attention available to patients. These wider health-system issues are important when judging the effectiveness of nursing care because even skilled nursing interventions may have limited impact when appropriate services are not available.

Need for a Person-Centred Approach

The overall analysis suggests that schizophrenia care should not be limited to controlling psychiatric symptoms. Reduction in hallucinations, delusions, and other acute symptoms is an important part of treatment, but it does not necessarily mean that the person has achieved recovery. Some individuals may continue to experience social isolation, lack of motivation, difficulty with self-care, unemployment, or low confidence.

A person-centred approach gives attention to the individual rather than focusing only on the diagnosis. Nurses should consider the patient's goals, preferences, strengths, abilities, and social situation when planning care. Patients should be encouraged to participate in decisions about their treatment as far as their condition allows. Setting small and realistic goals can help patients gradually improve their daily functioning and confidence. This approach also supports dignity, autonomy, and a greater sense of control over one's recovery.

Future Directions in Mental Health Nursing

Future mental health nursing should place greater emphasis on community-based and recovery-oriented care. Support should continue after hospital discharge rather than ending when acute symptoms improve. Regular follow-up can help nurses identify early difficulties, encourage treatment adherence, provide health education, support families, and connect patients with rehabilitation and community services.

Tele-mental health and digital technologies may provide additional support for continuing care. They can assist with follow-up, counselling, education, and communication with healthcare professionals, particularly for people who face difficulties travelling to specialist services. However, technology should be considered a supportive tool rather than a complete replacement for face-to-face nursing care. Some patients require direct assessment, personal interaction, and intensive support that cannot be fully provided through digital platforms.

There is also a need for more individualised and culturally appropriate nursing care. Patients differ in their symptoms, family situations, social environments, treatment preferences, and recovery goals. Future research should therefore examine which combinations of nursing interventions work best for different groups and healthcare settings. Research should also consider wider outcomes such as treatment adherence, quality of life, social participation, independence, employment, family wellbeing, and patient satisfaction rather than measuring success only through symptom reduction.

Critical Perspective

Overall, mental health nursing interventions can make a meaningful contribution to the management of schizophrenia, but their success depends on several factors beyond the intervention itself. Effective care requires adequately trained nurses, sufficient time, teamwork among healthcare professionals, family and community support, accessible mental health services, and active participation of patients in their care.

The strongest approach combines clinical management, psychosocial support, rehabilitation, and recovery-focused nursing care. Future practice should move towards an integrated, person-centred, recovery-oriented, and community-linked model. Such care recognises that successful management of schizophrenia involves more than controlling psychiatric symptoms. It also means helping individuals maintain relationships, manage everyday

activities, develop greater independence, participate in community life, and achieve a better quality of life.

Conclusion

Schizophrenia is a persistent mental health condition that can influence many areas of a person's life, including thinking, emotions, behaviour, communication, relationships, self-care, and independent living. Because of its wide-ranging effects, treatment cannot be limited to medication or control of acute psychiatric symptoms. Mental health nurses have an important role in providing continuous care and supporting patients at different stages of the illness. Assessment, therapeutic communication, medication-related support, patient education, safety management, family guidance, relapse prevention, rehabilitation, and follow-up can work together to support better long-term outcomes.

The review indicates that the success of nursing care varies from one individual to another. The patient's symptoms, cognitive abilities, personal preferences, family environment, social circumstances, and availability of healthcare services can all influence the outcome. In addition, problems such as irregular treatment use, stigma, cognitive limitations, caregiver stress, inadequate rehabilitation services, and shortage of trained mental health professionals may make effective care more difficult. Addressing these concerns requires cooperation among patients, families, nurses, mental health professionals, and community-based services.

An important finding of the analysis is that recovery in schizophrenia should not be judged only by whether psychiatric symptoms have decreased. Although symptom control remains necessary, meaningful improvement also includes better self-care, stronger social relationships, and participation in treatment, greater independence, community involvement, and improved quality of life. Therefore, nursing practice should move beyond a narrow symptom-based model and give greater importance to person-centred and recovery-oriented care. Nurses can support this process by understanding individual needs, recognising personal strengths, encouraging participation in care decisions, and setting achievable goals with patients.

Future mental health nursing should strengthen community follow-up, rehabilitation, family support, and continuity of care. Tele-mental health and other digital approaches may provide additional support where appropriate, particularly for follow-up and communication. At the same time, nursing interventions should be flexible enough to reflect individual, social, and cultural differences. Further research is required to identify effective combinations of nursing interventions and to examine their effects on outcomes such as quality of life, independence, social functioning, treatment participation, and family wellbeing.

In conclusion, mental health nursing is an essential part of comprehensive schizophrenia care. The purpose of nursing intervention should extend beyond managing symptoms to helping individuals live more stable, independent, and meaningful lives. Effective schizophrenia care should promote not only clinical stability but also dignity, functional ability, social participation, and quality of life. An integrated model that brings together clinical treatment, psychosocial rehabilitation, family involvement, and community support can provide a stronger basis for sustained recovery.

References

1. World Health Organization. Schizophrenia. World Health Organization,2025. https://www.who.int/news-room/fact-sheets/schizophrenia
2. World Health Organization. Guidance on Community Mental Health Services: Promoting Person-Centred and Rights-Based Approaches. World Health Organization,2021. https://www.who.int/publications/i/item/9789240025707
3. Directorate General of Health Services, Ministry of Health and Family Welfare, Government of India. National Mental Health Programme. Government of India,2026. https://dghs.mohfw.gov.in/national-mental-health-programme.php
4. Ministry of Health and Family Welfare, Government of India. Operational Guidelines: The National Tele Mental Health Programme of India—Tele MANAS. Ministry of Health and Family Welfare,2026:8. https://www.mohfw.gov.in/sites/default/files/Operational%20Guidelines%20-%20Tele%20Manas.pdf
5. Ministry of Health and Family Welfare, Government of India. National Mental Health Policy 2014. Ministry of Health and Family Welfare,2014:4–5. https://www.nhm.gov.in/images/pdf/National_Health_Mental_Policy.pdf
6. Harris BA, Panozzo G. Therapeutic alliance, relationship building, and communication strategies for the schizophrenia population: An integrative review. Archives of Psychiatric Nursing,2019:33(1):104–111. https://doi.org/10.1016/j.apnu.2018.08.003
7. Issac A, *et al*. Barriers and facilitators to medication adherence among individuals with mental disorders: A qualitative systematic review and evidence synthesis. Nursing & Health Sciences,2025:27(2):e70153. https://doi.org/10.1111/nhs.70153
8. Thongsalab J, Yunibhand J, Uthis P. Recovery-oriented nursing service for people with schizophrenia in the community: An integrative review. Belitung Nursing Journal,2023:9(3):198–208. https://doi.org/10.33546/bnj.2632
9. Harris BA, Panozzo G. Therapeutic alliance, relationship building, and communication strategies for the schizophrenia population: An integrative review. Archives of Psychiatric Nursing,2019:33(1):104–111. https://doi.org/10.1016/j.apnu.2018.08.003
10. Issac A, *et al*. Barriers and facilitators to medication adherence among individuals with mental disorders: A qualitative systematic review and evidence synthesis. Nursing & Health Sciences,2025:27(2):e70153. https://doi.org/10.1111/nhs.70153
11. Price D, Baker S. Key components of de-escalation techniques: A thematic synthesis. International Journal of Mental Health Nursing,2012:21(4):310–319. https://doi.org/10.1111/j.1447-0349.2011.00893.x
12. Zhou MH, *et al*. Psycho education for schizophrenia: A systematic review and meta-analysis. Psychiatry Research,2020:293:113453. https://doi.org/10.1016/j.psychres.2020.113453
13. Pharoah EM, *et al*. Family intervention for schizophrenia. Cochrane Database of Systematic Reviews,2010(12):CD000088. https://doi.org/10.1002/14651858.CD000088.pub3

14. Birchwood M, *et al*. Early intervention in psychosis: The critical period hypothesis. British Journal of Psychiatry,1998:172(S33):53–59. https://doi.org/10.1192/bjp.172.33.53
15. Corrigan PW, Drake RE, Rüsch MAR. The effects of illness management and recovery on self-determination and quality of life in individuals with serious mental illness. Psychiatric Services,2010:61(10):1026–1030. https://doi.org/10.1176/ps.2010.61.10.1026
16. Priebe S, *et al*. Effectiveness of community mental health team management in severe mental illness: A systematic review and meta-analysis. The Lancet,2006:367(9516):1841–1850. https://doi.org/10.1016/S0140-6736(06)68802-7
17. Hormazábal-Salgado R, Whitehead D, Osman AD, Hills D. Person-centred decision-making in mental health: A scoping review. Issues in Mental Health Nursing,2024:45(3):294–310. https://doi.org/10.1080/01612840.2023.2288181