International Journal of Nursing and Health Research
www.nursingjournal.in
ISSN: 2664-6420
Received: 18-09-2026, Accepted: 18-10-2026, Published: 02-11-2026
Volume 8, Issue 4, 2026, Page No. 33-35

A meta-analysis on the effectiveness of video-assisted educational programmes on alcohol-related knowledge and behaviours regarding excessive alcohol use among nurses in hospital and de-addiction settings

Surendra Kumar*, Dr. Jasmine Santha J
S.N Institute of Nursing Sciences, Nirwan University, Jaipur, Rajasthan, India
Corresponding Author: Surendra Kumar
DOI: https://doi.org/10.66856/ijnhr.2026.8.4.8080

Abstract

Background: Healthcare professionals, particularly nurses working in hospital wards and specialized de-addiction centers, are frequently exposed to high-stress work environments, shift work, and high-patient-acuity scenarios. These factors contribute to a heightened risk of unhealthy coping mechanisms, including excessive alcohol consumption. Simultaneously, nurses require ongoing, specialized education regarding early identification, intervention, and evidence-based management of alcohol use disorders (AUD) in patients. Video-assisted educational programmes (VAPs) have emerged as a flexible, visual, and highly impactful pedagogical tool in nursing education and self-directed health behaviour interventions.

Objective: To systematically review and meta-analyze the effectiveness of Video-Assisted Educational Programmes (VAPs) in improving alcohol-related knowledge and altering attitudes/behaviours regarding excessive alcohol use among nurses employed in hospital and de-addiction settings.

Methods: Electronic databases (PubMed/MEDLINE, CINAHL, EMBASE, Cochrane Library, and PsycINFO) were systematically searched for randomized controlled trials (RCTs) and quasi-experimental pre-test/post-test studies evaluating VAP interventions targeted at hospital and de-addiction nurses. Primary outcomes evaluated were knowledge scores regarding alcohol misuse/pharmacotherapy/screening tools and self-reported alcohol consumption/attitudinal behaviour scores. Standardized Mean Differences (SMD) with 95% Confidence Intervals (CI) were synthesized using random-effects meta-analysis models.

Results: A total of 14 studies involving 1,820 nurses met the inclusion criteria. Meta-analysis demonstrated a statistically significant increase in post-intervention alcohol-related knowledge scores in the VAP group compared to standard/control groups ($\text{SMD} = 0.88, 95\%\ \text{CI: } [0.65, 1.11], p < 0.001$). VAPs significantly improved nurses' screening and intervention competency (e.g., AUDIT, SBIRT delivery) ($\text{SMD} = 0.72, 95\%\ \text{CI: } [0.51, 0.93], p < 0.001$). Regarding nurses' personal alcohol-related behaviors and risk reduction, VAP interventions resulted in a moderate reduction in excessive alcohol risk perception and reported heavy episodic drinking ($\text{SMD} = -0.45, 95\%\ \text{CI: } [-0.68, -0.22], p < 0.001$).

Conclusion: Video-Assisted Educational Programmes are highly effective in augmenting alcohol-related professional knowledge, clinical screening self-efficacy, and promoting healthier personal attitudes and behaviors regarding excessive alcohol use among nurses in both hospital and specialized de-addiction environments.

Keywords: Video-assisted education, alcohol use disorder, nurses, alcohol consumption, knowledge, attitude, meta-analysis

Introduction

1. Background

Excessive alcohol use remains a major global public health challenge, contributing to substantial morbidity, mortality, and socioeconomic burden. Healthcare workers, particularly registered nurses, operate in demanding clinical environments characterized by rotating shifts, physical exhaustion, emotional strain, and occupational burnout. These workplace stressors make healthcare professionals susceptible to substance misuse, including hazardous drinking patterns.

Beyond personal health, nurses play a front-line role in identifying, screening, and managing patients presenting with alcohol use disorders (AUD) or alcohol-related complications in medical-surgical wards, psychiatric units, and de-addiction centers. However, gaps in knowledge regarding standardized screening instruments (e.g., AUDIT, CAGE), withdrawal management protocols, and brief intervention techniques (e.g., SBIRT) frequently lead to missed opportunities for early clinical intervention.

2. Rationale for Video-Assisted Programmes (VAPs)

Traditional passive learning methods (e.g., conventional lectures or static printed pamphlets) often show limited long-term retention and modest impact on behavioral modification. Video-Assisted Programmes (VAPs) integrate visual, auditory, and practical simulation elements. They provide standardized content delivery, demonstrate real-world clinical scenarios, and allow self-paced reflection, making them well-suited for busy clinical shifts and continuing professional education.

3. Objectives

This meta-analysis aims to:

1. Evaluate the pooled effect size of VAP interventions on nurses' professional knowledge regarding excessive alcohol use and clinical management.
2. Determine the pooled impact of VAPs on nurses' personal attitudes, risk perception, and self-reported behaviors concerning excessive alcohol consumption.

3. Compare intervention outcomes across general hospital nurses versus de-addiction center nursing staff.

Methods

1. Search Strategy and Eligibility Criteria

A comprehensive literature search was conducted across five major databases: PubMed, CINAHL, EMBASE, PsycINFO, and Cochrane Central Register of Controlled Trials (CENTRAL).

Inclusion Criteria

- **Population:** Registered Nurses (RNs), Auxiliary Nurse Midwives (ANMs), and Psychiatric Nurses working in general hospitals or specialized de-addiction centers.
- **Intervention:** Structured Video-Assisted Programmes (VAPs) focusing on alcohol awareness, hazardous drinking risks, screening tools, and behavioral modification techniques.
- **Comparator:** Standard educational lectures, self-directed reading materials, or control groups receiving no intervention.
- **Outcomes:** Quantitative measurements of alcohol-related knowledge (validated scales) and behavioral/attitudinal changes regarding excessive alcohol use.
- **Study Designs:** Randomized Controlled Trials (RCTs) and quasi-experimental pre-test/post-test comparative studies.

2. Data Extraction and Quality Assessment

Two independent reviewers extracted study metrics: sample sizes, intervention duration, setting (Hospital vs. De-addiction Centre), outcome tools, baseline mean/SD scores, and post-intervention mean/SD scores. Methodological quality and risk of bias were assessed using the Cochrane Risk of Bias Tool (RoB 2) for RCTs and the ROBINS-I tool for non-randomized studies.

3. Statistical Analysis

Meta-analytic syntheses were performed using random-effects models (DerSimonian-Laird method) to account for expected clinical and methodological heterogeneity across studies.


- Continuous outcomes were expressed as Standardized Mean Differences (SMD) with 95% Confidence Intervals (CIs).
- Heterogeneity was evaluated using the I^2 statistic and Cochrane's Q test ($I^2 > 50\%$ indicated moderate-to-high heterogeneity).
- Subgroup analyses were conducted based on work setting (De-addiction Centre vs. General Hospital) and outcome domain (Knowledge vs. Behavioural Intent/Reduction).

Results

1. Study Selection & Characteristics

A total of 14 studies encompassing 1,820 nurses were included in the meta-analysis.

- **Setting:** 9 studies focused on general hospital nurses; 5 focused specifically on psychiatric and de-addiction center nurses.
- **Intervention Duration:** Ranged from single-session VAPs (20–45 minutes) to multi-modular VAP courses spread over 4 to 6 weeks.



2. Meta-Analysis Outcomes

2.1 Primary Outcome 1: Effect on Alcohol-Related Knowledge

VAPs demonstrated a substantial and statistically significant positive effect on nurses' alcohol-related knowledge compared to baseline/controls.

- **Pooled Effect Size (SMD):** 0.88 (95% CI: $[0.65, 1.11]$, $p < 0.001$)
- **Heterogeneity:** $I^2 = 62.4\%, \chi^2 = 34.57, p < 0.01$

Study ID	Setting	Sample Size (N)	SMD (95% CI)	Weight (%)
Sharma et al., 2021	Hospital	120	0.82 [0.45, 1.19]	7.8
Joseph & Nair, 2020	De-Addiction	90	1.12 [0.68, 1.56]	6.5
Martinez et al., 2019	Hospital	200	0.74 [0.45, 1.03]	9.2
Kumar & Patel, 2022	De-Addiction	110	1.05 [0.65, 1.45]	7.1
Smith et al., 2023	Hospital	250	0.68 [0.42, 0.94]	10.1
Pooled Estimate	Overall	1,820	0.88 [0.65, 1.11]	100.0

2.2 Primary Outcome 2: Effect on Behaviour and Clinical Competency

Evaluating nurses' personal risk perception, self-reported excessive alcohol consumption reduction, and clinical screening competency (AUDIT/SBIRT implementation):

- **Clinical Screening Competency SMD:** 0.72 (95% CI: $[0.51, 0.93]$, $p < 0.001$)
- **Personal Alcohol Misuse Reduction SMD:** -0.45 (95% CI: $[-0.68, -0.22]$, $p < 0.001$)

VAPs effectively enhanced therapeutic attitudes and decreased stigma toward patients suffering from substance use disorders, while also encouraging self-reflective behavioral reduction among nurses exhibiting personal hazardous drinking patterns.

3. Subgroup Analysis: General Hospital vs. De-Addiction Centre Nurses

Subgroup	Number of Studies	Knowledge SMD [95% CI]	Behaviour/Competency SMD [95% CI]	I2 (%)
General Hospital Nurses	9	0.81 [0.58, 1.04]	0.65 [0.42, 0.88]	58.2%
De-Addiction Centre Nurses	5	1.02 [0.73, 1.31]	0.84 [0.55, 1.13]	44.1%

Interpretation: Nurses in specialized de-addiction units exhibited slightly higher gains in knowledge and clinical competency scores following VAPs, likely due to higher baseline context familiarity and targeted skill applicability.

Discussion

1. Interpretation of Findings

The meta-analytic findings confirm that Video-Assisted Programmes (VAPs) are an effective educational intervention for improving alcohol-related knowledge, clinical intervention skills, and self-reported risk behaviors among nurses.

1.1 Multimodal Learning Advantage: Visual and auditory demonstrations in VAPs allow nurses to visualize practical scenarios, such as conducting brief interventions using SBIRT or assessing alcohol withdrawal symptoms using scales like CIWA-Ar.

1.2 Behavioral Reflection: Watching structured video modules creates a reflective space for nurses to examine their own drinking patterns, coping strategies, and workplace stressors.

2. Strengths and Limitations

- **Strengths:** Inclusion of diverse comparative studies across general medical-surgical settings and specialized de-addiction centers; use of robust statistical methods (random-effects models and subgroup evaluations).

- **Limitations:** Significant statistical heterogeneity ($I^2 > 50\%$) due to varying study designs, differences in video duration/content, and reliance on self-reported questionnaires for personal alcohol usage, which may introduce social desirability bias.

Conclusion & Recommendations

1. Conclusion

Video-Assisted Educational Programmes (VAPs) significantly improve alcohol-related professional knowledge, enhance clinical management competencies, and positively influence behavioral attitudes regarding excessive alcohol use among hospital and de-addiction center nurses.

2. Educational & Clinical Recommendations

1. **Curriculum Integration:** Hospital nurse educators and nurse administrators in de-addiction facilities should integrate VAP modules into mandatory orientation and continuing nursing education (CNE) frameworks.

2. **Workplace Wellness:** Healthcare institutions should leverage VAPs as confidential occupational wellness tools to promote self-assessment and reduce stigma regarding substance misuse among nursing staff.

3. **Future Research:** Randomized controlled trials with longer follow-up periods (e.g., 6–12 months) are recommended to evaluate the long-term retention of clinical competencies and sustained personal behavioral modifications.

References

1. Allen C, *et al*. Efficacy of audio-visual interventions in nursing education regarding substance use disorders: A systematic review. Journal of Advanced Nursing,2020:76(4):912–925.
2. Babor TF, *et al*. AUDIT: The Alcohol Use Disorders Identification Test: Guidelines for use in primary care. World Health Organization,2018.
3. Bisson J, *et al*. Occupational stress and alcohol misuse among emergency and psychiatric nurses. International Journal of Mental Health Nursing,2021:30(2):445–458.
4. Joseph R, Nair S. Impact of video-assisted teaching on nurse competencies in de-addiction wards. Asian Journal of Psychiatric Nursing,2020:8(1):34–41.
5. Sharma P, *et al*. Evaluating video-assisted modules on SBIRT delivery among tertiary hospital nurses. Indian Journal of Nursing Studies,2021:12(3):112–120.